



# Human IL-1 $\beta$ premium grade

10  $\mu$ g  
25  $\mu$ g  
100  $\mu$ g  
1000  $\mu$ g

130-093-897  
130-093-563  
130-093-898  
130-093-899

## Contents

1. Description
  - 1.1 Background information
  - 1.2 Applications
2. References

## 1. Description

<b>Components</b>	<b>Human IL-1<math>\beta</math>, premium grade:</b> Purified recombinant human interleukin 1 $\beta$ (153 amino acid residues).
<b>Sizes</b>	10 $\mu$ g, 25 $\mu$ g, 100 $\mu$ g, 1000 $\mu$ g.
<b>Biological activity</b>	The ED <sub>50</sub> is <0.03 ng/mL* corresponding to a specific activity of >3 $\times$ 10 <sup>7</sup> I.U./mg.
<b>Molecular mass</b>	17.4 kDa.
<b>Source</b>	Produced in <i>E. coli</i> .
<b>Product format</b>	Lyophilized from a 0.2 $\mu$ m filtered solution.
<b>Stabilizer</b>	None (or trehalose and mannitol, 25 $\mu$ g size).
<b>Purity</b>	>97% as determined by gel filtration or SDS PAGE analysis.
<b>Endotoxin level</b>	Low endotoxin (<1 EU/ $\mu$ g cytokine) as determined by Limulus Amebocyte Lysate (LAL) assay.
<b>Storage</b>	Lyophilized Human IL-1 $\beta$ , premium grade should be stored at -20 °C. The expiration date is indicated on the vial label. Upon reconstitution aliquots should be stored at -20 °C. Avoid repeated freeze-thaw cycles.
<b>Reconstitution</b>	It is recommended to reconstitute lyophilized Human IL-1 $\beta$ with deionized sterile-filtered water up to a final concentration of 10 $\mu$ g/mL. <b>▲ Note:</b> Addition of carrier protein, such as 0.1% bovine serum albumin (BSA) or human serum albumin (HSA) may have stabilizing effects. Further dilutions should be prepared with 1% BSA or HSA in phosphate-buffered saline (PBS).

\* The ED<sub>50</sub> is determined by proliferation assay using D10.G4.1 or TF-1 cells according to Poole and Gaines-Das<sup>1</sup> and Kitamura *et al.*<sup>2</sup>, respectively. The proliferation assay was calibrated with the first international reference standard for human IL-1 $\beta$  (NIBSC code 86/680) provided by the WHO/National Institute for Biological Standards and Control.

## 1.1 Background information

Interleukin 1 $\beta$  (IL-1 $\beta$ ) is a proinflammatory cytokine that is secreted mainly by monocytes and macrophages. IL-1 $\beta$  secretion has also been reported for a variety of other cells, including B cells, NK cells, dendritic cells, astrocytes, and microglial cells. It mediates inflammatory responses in B cells, T cells, and NK cells by inducing the production of cytokines, such as IL-2, IL-3, IL-6, as well as interferons. Upon exposure to IL-1 $\beta$ , endothelial cells and smooth muscle cells synthesize prostaglandins and other derivatives of arachidonic acid. In addition, IL-1 $\beta$  is found in synovial fluid of arthritis patients, causing degranulation of basophils and eosinophils as well as activation of osteoclasts. IL-1 $\beta$  is mitogenic for mesangial cells, glial cells, and keratinocytes. It has been shown that IL-1 $\beta$  is a potent modulator of CD40L-induced cytokine secretion by human dendritic cell (DC) subsets, such as monocyte-derived dendritic cells (MoDCs), CD34<sup>+</sup>-derived DCs, and peripheral blood DCs.<sup>3</sup>

## 1.2 Applications

- IL-1 $\beta$  can be used for a variety of applications, including induction of MoDC maturation, chemotaxis assays, and investigation of IL-1 receptor signaling.

Optimal concentration for a specific application should be determined by a dose-response experiment.

## 2. References

1. Poole, S. and Gaines-Das, R. E. (1991) The international standards for interleukin-1 alpha and interleukin-1 beta. Evaluation in an international collaborative study. *J. Immunol. Methods* 142: 1-13.
2. Kitamura, T. *et al.* (1991) IL-1 up-regulates the expression of cytokine receptors on a factor-dependent human hemopoietic cell line, TF-1. *Int. Immunol.* 3: 571-577.
3. Luft, T. *et al.* (2002) IL-1 beta enhances CD40 ligand-mediated cytokine secretion by human dendritic cells (DC): a mechanism for T cell-independent DC activation. *J. Immunol.* 168: 713-722.

All protocols and data sheets are available at [www.miltenyibiotec.com](http://www.miltenyibiotec.com).

## Warranty

The products sold hereunder are warranted only to be free from defects in workmanship and material at the time of delivery to the customer. Miltenyi Biotec GmbH makes no warranty or representation, either expressed or implied, with respect to the fitness of a product for a particular purpose. There are no warranties, expressed or implied, which extend beyond the technical specifications of the products. Miltenyi Biotec GmbH's liability is limited to either replacement of the products or refund of the purchase price. Miltenyi Biotec GmbH is not liable for any property damage, personal injury or economic loss caused by the product.

Unless otherwise specifically indicated, Miltenyi Biotec products and services are for research use only and not for diagnostic or therapeutic use.

MACS is a registered trademark of Miltenyi Biotec GmbH.

Copyright © 2008 Miltenyi Biotec GmbH. All rights reserved.

