



# Anti-IL-5 antibodies human

Anti-IL-5-PE  
Anti-IL-5-APC

130-091-648  
130-091-834

## Contents

1. Description
  - 1.1 Background information
  - 1.2 Applications
  - 1.3 Recommended antibody dilution
  - 1.4 Reagent requirements
2. Protocols
  - 2.1 Sample preparation
  - 2.2 *In vitro* stimulation of PBMCs
  - 2.3 Intracellular immunofluorescent cytokine staining
    - 2.3.1 Intracellular staining of cells in suspension
    - 2.3.2 Intracellular staining in combination with magnetic cell separation (solid phase intracellular staining)
3. Examples of intracellular cytokine staining with Anti-IL-5 antibodies
4. Appendix: Flask and dish sizes for *in vitro* stimulation

## 1. Description

<b>Components</b>	1 mL Anti-IL-5 antibodies, human: monoclonal Anti-IL-5 antibodies conjugated to R-phycoerythrin (PE), or allophycocyanin (APC).
<b>Clone</b>	JES1-39D10 (isotype: rat IgG2a).
<b>Capacity</b>	100 tests or up to 10 <sup>9</sup> total cells.
<b>Product format</b>	Antibodies are supplied in buffer containing stabilizer and 0.05% sodium azide.
<b>Storage</b>	Store protected from light at 2–8 °C. Do not freeze. The expiration date is indicated on the vial label.

### 1.1 Background information

Interleukin 5 (IL-5) is a cytokine predominantly secreted by CD4<sup>+</sup> T cells. It is involved in a range of allergic reactions and mediates immune reactions against parasites. IL-5 also acts on other cell types, like B cells.

Anti-IL-5 antibodies are designed for intracellular staining of IL-5-producing cells. Cells can be stimulated for IL-5-production, for example, by polyclonal stimulation with mitogens. For induction of IL-5 production by antigen-specific T cells, cells are restimulated with respective antigen. IL-5 can be accumulated in the cells by addition of secretion inhibitors like brefeldin A. After fixation and permeabilization of the cell sample, IL-5-producing cells can be stained intracellularly with Anti-IL-5 antibodies. Staining of surface markers allows simultaneous flow cytometric analysis of subsets and activation status of the IL-5-producing cells.

Magnetically enriched cells can be stained intracellularly for IL-5 production directly on the MACS® Column. This procedure ensures higher sensitivity of detection and minimizes loss of cells during washing procedures. The protocol is useful for cytokine analysis of rare cells, for example, CD4<sup>+</sup> T cells in HIV patients, or other cell sources than PBMCs, for example, bronchoalveolar lavages, or synovial fluids. For details see section 2.3.2.

### 1.2 Applications

- Flow cytometric identification and enumeration of IL-5-producing cells upon polyclonal stimulation with mitogens.
- Identification and enumeration of IL-5-producing antigen-specific T cells upon restimulation with the respective antigen.
- Monitoring of specificity of antigen-specific T cell lines.
- Analysis of cytokine production of rare cells by solid-phase staining technology, in combination with magnetic enrichment of cells. For details see section 2.3.2.

### 1.3 Recommended antibody dilution

- Anti-IL-5 antibodies should be used at a dilution of 1:10.
- Cross-reactivity: The monoclonal Anti-IL-5 antibody is reported to react with IL-5 produced by a subset of peripheral blood mononuclear cells (PBMCs) from rhesus monkey (*Macaca mulatta*) and cynomolgus monkey (*Macaca fascicularis*).

### 1.4 Reagent requirements

- Buffer: Prepare a solution containing phosphate-buffered saline (PBS), pH 7.2, 0.5% bovine serum albumin (BSA), and 2 mM EDTA by diluting MACS BSA Stock Solution (# 130-091-376) 1:20 with autoMACS™ Rinsing Solution (# 130-091-222). Keep buffer cold (2–8 °C).  
▲ **Note:** EDTA can be replaced by other supplements such as anticoagulant citrate dextrose formula-A (ACD-A) or citrate phosphate dextrose (CPD). BSA can be replaced by other proteins such as human serum albumin, human serum, or fetal bovine serum. Buffers or media containing Ca<sup>2+</sup> or Mg<sup>2+</sup> are not recommended for use.
- Culture medium, for example, RPMI 1640 (# 130-091-440) containing 5% human serum, e.g., autologous or AB serum (do not use BSA or FCS because of non-specific stimulation!).
- Reagents for T cell stimulation, such as phorbol myristate acetate (PMA)/ionomycin, antigenic peptide or protein, e.g., peptide pools (PepTivators), or CMV pp65 – Recombinant Protein (# 130-091-824). For details see the respective data sheet. For more information about antigens see [www.miltenyibiotec.com](http://www.miltenyibiotec.com).
- Secretion inhibitor, e.g., brefeldin A.
- Inside Stain Kit (# 130-090-477) for fixation and permeabilization of cells.



- Fluorochrome-conjugated antibodies for surface staining, e.g., CD4-PE (# 130-091-231), and/or CD8-FITC (# 130-080-601) or CD69-FITC (# 130-092-166).

### Additional requirements for intracellular cytokine staining in combination with magnetic cell separation (see protocol 2.3.2)

- MACS MicroBeads, e.g., CD4 MicroBeads (# 130-045-101).
- MS Columns and suitable MACS Separator (MiniMACS™, OctoMACS™, VarioMACS™, or SuperMACS™ Separator).
  - ▲ **Note:** Column adapters are required to insert certain columns into VarioMACS or SuperMACS separators. For details see the respective MACS Separator data sheet.
- (Optional) Pre-Separation Filters (# 130-041-407) to remove cell clumps.

## 2. Protocols

### 2.1 Sample preparation

When working with anticoagulated peripheral blood or buffy coat, peripheral blood mononuclear cells (PBMCs) should be isolated by density gradient centrifugation, for example using Ficoll-Paque™. For details see the General Protocols section of the respective separator user manual. The General Protocols are also available at [www.miltenyibiotec.com/protocols](http://www.miltenyibiotec.com/protocols).

▲ **Note:** To remove platelets after density gradient separation, resuspend cell pellet in buffer and centrifuge at 200×g for 10–15 minutes at 20 °C. Carefully aspirate supernatant. Repeat washing step.

When working with tissues or lysed blood, prepare a single-cell suspension using standard methods. For details see the General Protocols section of the respective separator user manual. General Protocols are also available at [www.miltenyibiotec.com/protocols](http://www.miltenyibiotec.com/protocols).

### 2.2 *In vitro* stimulation of PBMCs

▲ Always include a negative control in the experiment. The sample should be treated exactly the same way as the stimulated sample, except for the addition of the stimulus.

▲ A positive control may also be included in the experiment, such as a sample stimulated with PMA/ionomycin as high control or SEB as medium control for IL-5-producing cells.

▲ Do not use media containing any non-human proteins, such as BSA or FCS, because of non-specific stimulation.

1. Wash cells by adding medium and centrifuge at 300×g for 10 minutes. Aspirate supernatant.
2. Resuspend cells at a density of 10<sup>7</sup> per mL in culture medium containing 5% human serum. Plate cells in dishes at a density of 5×10<sup>6</sup> cells/cm<sup>2</sup>. For details see 4. Appendix: Flask and dish sizes for *in vitro* stimulation.
3. Add antigen or control reagent:
  - 1–10 µg/mL peptide
  - 10–100 µg/mL protein
  - 1 µg/mL SEB
  - 20 ng/mL PMA and 1 µg/mL ionomycin
4. Incubate cells for 2 hours at 37 °C and 5–7% CO<sub>2</sub>.
5. Add 1 µg/mL brefeldin A and incubate for an additional 4 hours at 37 °C and 5–7% CO<sub>2</sub>.

6. Collect cells carefully by pipetting up and down when working with smaller volumes, or by using a cell scraper. Rinse the dish with cold buffer. Check microscopically for any remaining cells, if necessary, rinse the dish again.

### 2.3 Intracellular immunofluorescent cytokine staining

#### 2.3.1 Intracellular staining of cells in suspension

▲ It is recommended to stain 10<sup>6</sup> cells per sample. When working with up to 10<sup>7</sup> cells, use the same volumes as indicated. When working with higher cell numbers, scale up all reagent volumes and total volumes accordingly (e.g., for 2×10<sup>7</sup> nucleated cells, use twice the volume of all indicated reagent volumes and total volumes).

1. Wash up to 10<sup>7</sup> cells by adding 1–2 mL of buffer and centrifuge at 300×g for 10 minutes. Aspirate supernatant completely.
2. (Optional) Stain cell surface antigens that are sensitive to fixation according to the manufacturer's recommendations. Then wash cells by adding 1–2 mL of buffer and centrifuge at 300×g for 10 minutes. Aspirate supernatant completely.
3. Resuspend up to 10<sup>7</sup> cells in 500 µL of buffer.
4. Add 500 µL of Inside Fix (Inside Stain Kit). Mix well and incubate for 20 minutes at room temperature.
5. Centrifuge at 300×g for 5 minutes. Aspirate supernatant carefully.
6. Wash cells by adding 1 mL of buffer and centrifuge at 300×g for 5 minutes. Aspirate supernatant carefully.
  - ▲ **Note:** Fixed cells may be stored at 2–8 °C for up to 1 week.
7. (Optional) Stain cell surface antigens with antibodies that are sensitive to permeabilization according to the manufacturer's recommendations, e.g., add 10 µL of CD4-PE (# 130-091-231), mix well and incubate for 10 minutes at 2–8 °C. Then wash cells by adding 1–2 mL of buffer and centrifuge at 300×g for 10 minutes. Aspirate supernatant completely.
8. Wash cells by adding 1 mL of Inside Perm (Inside Stain Kit). Centrifuge at 300×g for 5 minutes. Aspirate supernatant carefully.
9. Resuspend cells in 90 µL of Inside Perm. Add 10 µL of the Anti-IL-5 antibody.
10. (Optional) Add additional staining antibodies to the solution, for example, for the staining of cell surface antigens internalized upon cell activation, such as CD3 and TCR α/β, or for the staining of antigens accumulating in the cell, such as CD69.
11. Mix well and incubate for 10 minutes in the dark at room temperature.
12. Wash cells by adding 1 mL of Inside Perm and centrifuge at 300×g for 5 minutes. Aspirate supernatant carefully.
13. Resuspend cell pellet in a suitable amount of buffer for analysis by flow cytometry or fluorescence microscopy. Store cells at 2–8 °C in the dark until analysis. Mix well before flow cytometric data acquisition.
  - ▲ **Note:** Samples may be stored at 2–8 °C in the dark for up to 24 hours.
  - ▲ **Note:** Do not use propidium iodide (PI) or 7-AAD staining.

### 2.3.2 Intracellular staining in combination with magnetic cell separation (solid phase intracellular staining)

▲ Work fast, keep cells cold, and use pre-cooled solutions. This will prevent capping of antibodies on the cell surface and non-specific cell labeling.

▲ Volumes for magnetic labeling given below are for up to  $10^7$  total cells. When working with fewer than  $10^7$  cells, use the same volumes as indicated. When working with higher cell numbers, scale up all reagent volumes and total volumes accordingly (e.g. for  $2 \times 10^7$  total cells, use twice the volume of all indicated reagent volumes and total volumes).


▲ For optimal performance it is important to obtain a single-cell suspension before magnetic separation. Pass cells through 30  $\mu$ m nylon mesh (Pre-Separation Filters, # 130-041-407) to remove cell clumps which may clog the column. Wet filter with buffer before use.

1. Wash cells by adding 1–2 mL of buffer per  $10^7$  cells and centrifuge cell suspension at  $300 \times g$  for 10 minutes. Aspirate supernatant completely.
2. Resuspend cell pellet in 80  $\mu$ L of buffer per  $10^7$  total cells.
3. Add 20  $\mu$ L of MACS MicroBeads, e.g., CD4 MicroBeads (# 130-045-101) per  $10^7$  total cells.  
▲ **Note:** For details on the procedure refer to the respective MACS MicroBead data sheet.
4. Mix well and incubate for 15 minutes in the refrigerator (2–8 °C).  
▲ **Note:** Working on ice may require increased incubation times. Higher temperatures and/or longer incubation times may lead to non-specific cell labeling.
5. (Optional) Stain cell surface antigens with antibodies that are sensitive to fixation according to the manufacturer's recommendations, e.g., add 10  $\mu$ L of CD4-PE (# 130-091-231), and incubate for additional 5 minutes at 2–8 °C. Then wash cells by adding 1–2 mL of buffer per  $10^7$  cells and centrifuge at  $300 \times g$  for 10 minutes. Aspirate supernatant completely.
6. Resuspend cell pellet in 500  $\mu$ L of buffer.
7. Place MS Column in the magnetic field of a suitable MACS Separator. For details see MS Column data sheet.
8. Prepare column by rinsing with 500  $\mu$ L of buffer.
9. Apply cell suspension onto the column.
10. Collect unlabeled cells that pass through and wash column with  $3 \times 500$   $\mu$ L of buffer. Collect total effluent; this is the unlabeled cell fraction. Perform washing steps by adding buffer three times. Only add new buffer when the column reservoir is empty.
11. Remove column from the separator and place it on a suitable collection tube.
12. Pipette 500  $\mu$ L of buffer onto the column. Immediately flush out the magnetically labeled cells by firmly pushing the plunger into the column.
13. Add 500  $\mu$ L of Inside Fix to the eluted cell fraction and incubate for 20 minutes at room temperature.
14. Place a second MS Column in the magnetic field of a suitable MACS Separator and prepare column by rinsing with 500  $\mu$ L of buffer.

15. Apply the fixed cell suspension onto the column.
16. Wash cells by rinsing the column with  $1 \times 500$   $\mu$ L of buffer. Then permeabilize cells by rinsing the column with  $2 \times 500$   $\mu$ L of Inside Perm.
17. Prepare a solution of 10  $\mu$ L of Anti-IL-5 antibodies and 90  $\mu$ L of Inside Perm.
18. (Optional) Add additional staining antibodies to the solution, for example, for the staining of cell surface antigens internalized upon cell activation, such as CD3 and TCR  $\alpha/\beta$  or for the staining of antigens accumulating in the cell, such as CD69.  
▲ **Note:** Do not exceed the total solution volume of 150  $\mu$ L.
19. Apply the solution onto the column and incubate for 10 minutes at room temperature.  
▲ **Note:** The MACS Column has a flow-stop mechanism that will retain the solution in the column.
20. Wash cells by rinsing the column with  $2 \times 500$   $\mu$ L of Inside Perm followed by  $1 \times 500$   $\mu$ L of buffer.
21. Remove column from the separator and place it on a suitable collection tube.
22. Pipette 500  $\mu$ L of buffer onto the column. Immediately flush out magnetically labeled cells by firmly pushing the plunger into the column.
23. Cells are now ready for analysis. Store cells at 2–8 °C in the dark until analysis. Mix well before flow cytometric data acquisition.  
▲ **Note:** Samples may be stored at 2–8 °C in the dark for up to 24 hours.  
▲ **Note:** Do not use propidium iodide (PI) or 7-AAD staining.

### 3. Examples of intracellular staining with Anti-IL-5 antibodies

Human PBMCs were incubated with and without 20 ng/mL PMA and 1 µg/mL ionomycin for 6 hours. After 2 hours brefeldin A was added. Cells were then fixed, permeabilized, and intracellularly stained with Anti-IL-5-PE or Anti-IL-5-APC. Cell surface staining was performed with CD4-PE (# 130-091-231) or CD4-APC (# 130-091-232) and CD69-FITC. Cells were analyzed by flow cytometry. Gating was performed according to the CD4 expression and side scatter properties of the cells. Autofluorescent cell debris was excluded in an FL-2 vs. FL-3 dot plot.



### 4. Appendix: Flask and dish sizes for *in vitro* stimulation

For *in vitro* stimulation of PBMCs (see 2.2) the cells should be resuspended in culture medium, containing 5% of human serum, at a dilution of  $10^7$  cells/mL. The cells should be plated at a density of  $5 \times 10^6$  cells/cm<sup>2</sup>. Both the dilution and the cell density are important to assure optimum stimulation.

The following table lists culture plate, dish and flask sizes suitable for different cell numbers. It also indicates the appropriate amount of medium to add.

Total cell number	Medium volume to add	Culture plate	Well diameter
$0.15 \times 10^7$	0.15 mL	96 well	0.64 cm
$0.50 \times 10^7$	0.50 mL	48 well	1.13 cm
$1.00 \times 10^7$	1.00 mL	24 well	1.60 cm
$2.00 \times 10^7$	2.00 mL	12 well	2.26 cm
$5.00 \times 10^7$	5.00 mL	6 well	3.50 cm
Total cell number	Medium volume to add	Culture dish	Dish diameter
$4.5 \times 10^7$	4.5 mL	small	3.5 cm
$10.0 \times 10^7$	10.0 mL	medium	6 cm
$25.0 \times 10^7$	25.0 mL	large	10 cm
$50.0 \times 10^7$	50.0 mL	extra large	15 cm
Total cell number	Medium volume to add	Culture flask	Growth area
$12 \times 10^7$	12 mL	50 mL	25 cm <sup>2</sup>
$40 \times 10^7$	40 mL	250 mL	75 cm <sup>2</sup>
$80 \times 10^7$	80 mL	720 mL	162 cm <sup>2</sup>
$120 \times 10^7$	120 mL	900 mL	225 cm <sup>2</sup>

All protocols and data sheets are available at [www.miltenyibiotec.com](http://www.miltenyibiotec.com).

#### Warnings

Reagents contain sodium azide. Under acidic conditions sodium azide yields hydrazoic acid, which is extremely toxic. Azide compounds should be diluted with running water before discarding. These precautions are recommended to avoid deposits in plumbing where explosive conditions may develop.

#### Warranty

The products sold hereunder are warranted only to be free from defects in workmanship and material at the time of delivery to the customer. Miltenyi Biotec GmbH makes no warranty or representation, either expressed or implied, with respect to the fitness of a product for a particular purpose. There are no warranties, expressed or implied, which extend beyond the technical specifications of the products. Miltenyi Biotec GmbH's liability is limited to either replacement of the products or refund of the purchase price. Miltenyi Biotec GmbH is not liable for any property damage, personal injury or economic loss caused by the product.

MACS is a registered trademark and autoMACS, MidiMACS, MiniMACS, OctoMACS, QuadroMACS, SuperMACS, and VarioMACS are trademarks of Miltenyi Biotec GmbH.

Ficoll-Paque is a trademark of GE Healthcare companies.

Copyright © 2008 Miltenyi Biotec GmbH. All rights reserved.