



August 2006

Cytokine Secretion Assays — reference list

Cytokine Secretion Assay technology

- Manz, R. *et al.* (1995) Analysis and Sorting of Live Cells According to Secreted Molecules Relocated to a Cell-Surface Affinity Matrix. *PNAS* 92: 1921–1925. [139]
- Brosterhus, H. *et al.* (1999) Enrichment and detection of live antigen-specific CD4⁺ and CD8⁺ T cells based on cytokine secretion. *Eur. J. Immunol.* 29: 4053–4059. [573]
- Desombre, I. *et al.* (2004) The interferon gamma secretion assay: a reliable tool to study interferon gamma production at the single cell level. *J. Immunol. Methods.* 286: 167–185. [4323]

Enumeration of cytokine-secreting cells

- Becker, C. *et al.* (2001) Adoptive tumor therapy with T lymphocytes enriched through an IFN- γ capture assay. *Nature Med.* 7: 1159–1162. [1207]
- Cohen, G.B. *et al.* (2002) Clonotype Tracking of TCR Repertoires during Chronic Virus Infections. *Virology* 304: 474–484. [2676]
- Akdis, M. *et al.* (2004) Immune Responses in Healthy and Allergic Individuals Are Characterized by a Fine Balance between Allergen-specific T Regulatory 1 and T Helper 2 Cells. *J. Exp. Med.* 199: 1567–1575. [3157]
- Desombre, I. *et al.* (2004) The interferon gamma secretion assay: a reliable tool to study interferon gamma production at the single cell level. *J. Immunol. Methods.* 286: 167–185. [4323]
- Veldman, C. *et al.* (2004) Type I Regulatory T Cells Specific for Desmoglein 3 Are More Frequently Detected in Healthy Individuals than in Patients with Pemphigus Vulgaris. *J. Immunol.* 172: 6468–6475. [5740]

Combination with peptide-MHC tetramer staining

- Pittet, M.J. *et al.* (2001) *Ex vivo* IFN- γ secretion by circulating CD8 T lymphocytes: Implications of a novel approach for T cell monitoring in infectious malignant diseases. *J. Immunol.* 166: 7634–7640. [1037]
- Meidenbauer, N. *et al.* (2003) Survival and Tumor Localization of Adoptively Transferred Melan-A-Specific T Cells in Melanoma Patients. *J. Immunol.* 170: 2161–2169. [2678]
- Morris, S.C. *et al.* (2006) IL-4 Induces *In Vivo* Production of IFN- γ by NK and NKT Cells. *J. Immunol.* 176: 5299–5305. [8659]

Adoptive transfer of cells isolated with the Cytokine Secretion Assay

- Oelke, M. *et al.* (2000) Generation and purification of CD8⁺ Melan-A-Specific Cytotoxic T Lymphocytes for Adoptive Transfer in Tumor Immunotherapy. *Clin. Cancer Res.* 6: 1997–2005. [663]
- Becker, C. *et al.* (2001) Adoptive tumor therapy with T lymphocytes enriched through an IFN- γ capture assay. *Nature Med.* 7: 1159–1162. [1207]
- Feuchtinger, T. *et al.* (2004) Isolation and expansion of human adenovirus-specific CD4⁺ and CD8⁺ T cells according to IFN-gamma secretion for adjuvant immunotherapy. *Exp. Hematol.* 32: 282–289. [4334]
- Rauser, G. *et al.* (2004) Rapid generation of combined CMV-specific CD4⁺ and CD8⁺ T-cell lines for adoptive transfer into recipients of allogeneic stem cell transplants. *Blood* 103: 3565–3572. [5341]
- Feuchtinger, T. *et al.* (2005) Detection of adenovirus-specific T cells in children with adenovirus infection after allogeneic stem cell transplantation. *Br. J. Haematol.* 128: 503–509. [5385]

Infectious diseases**CMV**

- Bissinger, A.L. *et al.* (2002) Isolation and expansion of human cytomegalo-virus-specific cytotoxic T lymphocytes using interferon- γ secretion assay. *Exp. Hematol.* 30: 1178–1184. [2158]
- Bitmansour, A.D. *et al.* (2002) Direct *Ex Vivo* Analysis of Human CD4⁺ Memory T Cell Activation Requirements at the Single Clonotype Level. *J. Immunol.* 169: 1207–1218. [2675]
- Rauser, G. *et al.* (2004) Rapid generation of combined CMV-specific CD4⁺ and CD8⁺ T-cell lines for adoptive transfer into recipients of allogeneic stem cell transplants. *Blood* 103: 3565–3572. [5341]
- Manley, T.J. *et al.* (2004) Immune evasion proteins of human cytomegalovirus do not prevent a diverse CD8⁺ cytotoxic T-cell response in natural infection. *Blood* 104: 1075–1082. [4345]

Hepatitis

- Meyer-Olson, D. *et al.* (2004) Limited T Cell Receptor Diversity of HCV-specific T Cell Responses Is Associated with CTL Escape. *J. Exp. Med.* 200: 307–319. [4343]
- Rico, M.A. *et al.* (2004) Virus-Specific Effector CD4⁺ T-Cell Responses in Hemodialysis Patients With Hepatitis C Virus Infection. *J. Med. Virol.* 72: 66–74. [4325]

HIV

- Altfeld, M. *et al.* (2001) HIV Study Collaboration. Vpr Is Preferentially Targeted by CTL During HIV-1 Infection. *J. Immunol.* 167: 2743–2752. [2495]
- Riberd, J.M. *et al.* (2001) Cutting Edge: Culture with High Doses of Viral Peptide Induces Previously Unprimed CD8⁺ T Cells to Produce Cytokine. *J. Immunol.* 167: 2437–2440. [2487]
- Yu, X.G. *et al.* (2002) Consistent Patterns in the Development and Immunodominance of Human Immunodeficiency Virus Type 1 (HIV-1)-Specific CD8⁺ T-Cell Responses following Acute HIV-1 Infection. *J. Virol.* 76: 8690–8701. [2681]
- Lee, S.K. *et al.* (2002) The functional CD8 T cell response to HIV becomes type-specific in progressive disease. *J. Clin. Invest.* 110: 1339–1347. [2677]
- Douek, D.C. *et al.* (2002) HIV preferentially infects HIV-specific CD4⁺ T cells. *Nature* 417: 95–98. [2320]
- Cohen, G.B. *et al.* (2002) Clonotype Tracking of TCR Repertoires during Chronic Virus Infections. *Virology* 304: 474–484. [2676]
- Lopes, A.R. *et al.* (2003) Greater CD8⁺ TCR Heterogeneity and Functional Flexibility in HIV-2 Compared to HIV-1 Infection. *J. Immunol.* 171: 307–316. [4698]
- Lichtenfeld, M. *et al.* (2004) HIV-1-specific cytotoxicity is preferentially mediated by a subset of CD8⁺ T cells producing both interferon- γ and tumor necrosis factor- α . *Blood* 104: 487–494. [4329]

EBV

- Bickham, K. *et al.* (2001) EBNA1-specific CD4⁺ T cells in healthy carriers of Epstein-Barr virus are primarily Th1 in function. *J. Clin. Invest.* 107: 121–130. [1035]
- Cohen, G.B. *et al.* (2002) Clonotype Tracking of TCR Repertoires during Chronic Virus Infections. *Virology* 304: 474–484. [2676]
- Koehne, G. *et al.* (2002) Quantitation, selection, and functional

characterization of Epstein-Barr virus-specific and alloreactive T cells detected by intracellular interferon- γ production and growth of cytotoxic precursors. *Blood* 99: 1730–1740. [2713]

- Paludan, C. *et al.* (2002) Epstein-Barr Nuclear Antigen 1-Specific CD4⁺ Th1 Cells Kill Burkitt's Lymphoma. *J. Immunol.* 169: 1593–1603. [2343]

Influenza

- Brosterhus, H. *et al.* (1999) Enrichment and detection of live antigen-specific CD4⁺ and CD8⁺ T cells based on cytokine secretion. *Eur. J. Immunol.* 29: 4053–4059. [573]
- Ponsaerts, P. *et al.* (2002) Messenger RNA Electroporation of Human Monocytes, Followed by Rapid *In Vitro* Differentiation, Leads to Highly Stimulatory Antigen-Loaded Mature Dendritic Cells. *J. Immunol.* 169: 1669–1675. [4331]
- Lucas, M. *et al.* (2004) *Ex Vivo* Phenotype and Frequency of Influenza Virus-specific CD4 Memory T Cells. *J. Virol.* 78: 7284–7287. [4314]
- Mayer, K.D. *et al.* (2005) The Functional Heterogeneity of Type 1 Effector T Cells in Response to Infection Is Related to the Potential for IFN- γ Production. *J. Immunol.* 174: 7732–7739. [7200]

Aspergillus

- Beck, O. *et al.* (2006) Generation of highly purified and functionally active human TH1 cells against *Aspergillus fumigatus*. *Blood* 107: 2562–2569. [5149]

Parasites

- Mohrs, K. *et al.* (2005) Systemic Dissemination and Persistence of Th2 and Type 2 Cells in Response to Infection with a Strictly Enteric Nematode Parasite. *J. Immunol.* 175: 5306–5313. [8122]

Tumor biology/ Tumor immunotherapy

- Oelke, M. *et al.* (2000) Generation and purification of CD8⁺ Melan-A-Specific Cytotoxic T Lymphocytes for Adoptive Transfer in Tumor Immunotherapy. *Clin. Cancer Res.* 6: 1997–2005. [663]
- Oelke, M. *et al.* (2000) Functional Characterization of CD8⁺ Antigen-Specific Cytotoxic T Lymphocytes after Enrichment Based on Cytokine Secretion: Comparison with the MHC-Tetramer Technology. *Scand. J. Immunol.* 52: 544–549. [970]
- Becker, C. *et al.* (2001) Adoptive tumor therapy with T lymphocytes enriched through an IFN- γ capture assay. *Nature Med.* 7: 1159–1162. [1207]
- Pittet, M.J. *et al.* (2001) *Ex vivo* IFN- γ secretion by circulating CD8 T lymphocytes: Implications of a novel approach for T cell monitoring in infectious malignant diseases. *J. Immunol.* 166: 7634–7640. [1037]
- Meyer zum Büschenfelde, K.H. *et al.* (2001) The Generation of Both T Killer and Th Cell Clones specific for the Tumor-Associated Antigen HER2 using retrovirally transduced Dendritic Cells. *J. Immunol.* 167: 1712–1719. [1260]
- Märten, A. *et al.* (2002) Generation of activated and antigen-specific T cells with cytotoxic activity after co-culture with dendritic cells. *Cancer Immunol. Immunother.* 51: 25–32. [2679]
- Valmori, D. *et al.* (2002) Circulating Tumor-reactive CD8⁺ T Cells in Melanoma Patients Contain a CD45RA⁺CCR7⁻ Effector Subset Exerting *ex Vivo* Tumor-specific Cytolytic Activity. *Cancer Res.* 62: 1743–1750. [6381]

- Rivoltini, L. *et al.* (2003) Human tumor-derived heat shock protein 96 mediates *in vitro* activation and *in vivo* expansion of melanoma- and colon carcinoma-specific T cells. *J. Immunol.* 171: 3467–3474. [4724]
- Consogno, G. *et al.* (2003) Identification of immunodominant regions among promiscuous HLA-DR-restricted CD4⁺ T-cell epitopes on the tumor antigen MAGE-3. *Blood* 101: 1038–1044. [4360]
- Meidenbauer, N. *et al.* (2003) Survival and Tumor Localization of Adoptively Transferred Melan-A-Specific T Cells in Melanoma Patients. *J. Immunol.* 169: 3652–3660. [2678]
- Ayyoub, M. *et al.* (2004) An immunodominant SSX-2–derived epitope recognized by CD4⁺ T cells in association with HLA-DR. *J. Clin. Invest.* 113: 1225–1233. [4340]
- Kloosterboer, F.M. *et al.* (2004) Direct Cloning of leukemia-reactive T cells from patients treated with donor lymphocyte infusions shows a relative dominance of hematopoiesis-restricted minor histocompatibility antigen HA-1 and HA-2 specific T cells. *Leukemia* 18: 798–808. [4322]
- Korangy, F. *et al.* (2004) Spontaneous tumor-specific humoral and cellular immune responses to NY-ESO-1 in hepatocellular carcinoma. *Clin. Cancer Res.* 10: 4332–4341. [9015]
- Sakai, Y. *et al.* (2004) Vaccination by genetically modified dendritic cells expressing a truncated neu oncogene prevents development of breast cancer in transgenic mice. *Cancer Research* 64: 8022–8028. [9017]
- Zhao, Y. *et al.* (2005) Primary Human Lymphocytes Transduced with NY-ESO-1 Antigen-Specific TCR Genes Recognize and Kill Diverse Human Tumor Cell Lines. *J. Immunol.* 174: 4415–4423. [6962]
- Zhang, Y. *et al.* (2005) Monitoring of Anti-Vaccine CD4 T Cell Frequencies in Melanoma Patients Vaccinated with a MAGE-3 Protein. *J. Immunol.* 174: 2404–2411. [6779]
- Goodyear, O. *et al.* (2005) CD8⁺ T cells specific for cancer germline gene antigens are found in many patients with multiple myeloma, and their frequency correlates with disease burden. *Blood* 106: 4217–4224. [8033]

Autoimmune disease

- Weissert, R. *et al.* (2002) High Immunogenicity of Intracellular Myelin Oligodendrocyte Glycoprotein Epitopes. *J. Immunol.* 169: 548–556. [2527]
- Veldman, C. *et al.* (2003) Dichotomy of Autoreactive Th1 and Th2 Cell Responses to Desmoglein 3 in Patients with Pemphigus Vulgaris (PV) and Healthy Carriers of PV-Associated HLA Class II Alleles. *J. Immunol.* 170: 635–642. [4327]
- Araki, M. *et al.* (2003) Th2 bias of CD4⁺ NKT cells derived from multiple sclerosis in remission. *Int. Immunol.* 15: 279–288. [4326]
- Veldman, C. *et al.* (2004) Type I Regulatory T Cells Specific for Desmoglein 3 Are More Frequently Detected in Healthy Individuals than in Patients with Pemphigus Vulgaris. *J. Immunol.* 172: 6468–6475. [5740]
- Veldman, C. *et al.* (2004) T Cell Recognition of Desmoglein 3 Peptides in Patients with Pemphigus Vulgaris and Healthy Individuals. *J. Immunol.* 172: 3883–3892. [3144]

Allergy

- Akdis, M. *et al.* (2000) Purification of allergen-specific T cells from allergic patients based on IL-4 and IFN- γ secretion. *MACS&more* 4/2: 9–10.
- Akdis, M. *et al.* (2004) Immune Responses in Healthy and Allergic Individuals Are Characterized by a Fine Balance between Allergen-specific T Regulatory 1 and T Helper 2 Cells. *J. Exp. Med.* 199: 1567–1575. [3157]
- Reefer, A.J. *et al.* (2004) A Role for IL-10-Mediated HLA-DR7-Restricted T Cell-Dependent Events in Development of the Modified Th2 Response to Cat Allergen. *J. Immunol.* 172: 2763–2772. [5536]
- Desombre, I. *et al.* (2004) The interferon gamma secretion assay: a reliable tool to study interferon gamma production at the single cell level. *J. Immunol. Methods.* 286: 167–185. [4323]

T cells

- Ouyang, W. *et al.* (2000) Stat6-Independent GATA-3 Autoactivation Directs IL-4-Independent Th2 Development and Commitment. *Immunity* 12: 27–37. [597]
- Novak, E.J. *et al.* (2001) Tetramer-Guided Epitope Mapping: Rapid Identification and Characterization of Immunodominant CD4⁺ T Cell Epitopes from Complex Antigens. *J. Immunol.* 166: 6665–6670. [2663]
- Smits, H.H. *et al.* (2001) IL-12-induced reversal of human Th2 cells is accompanied by full restoration of IL-12 responsiveness and loss of GATA-3 expression. *Eur. J. Immunol.* 31: 1055–1065. [983]
- Hornung, V. *et al.* (2002) Quantitative Expression of Toll-Like Receptor 1–10 mRNA in Cellular Subsets of Human Peripheral Blood Mononuclear Cells and Sensitivity to CpG Oligodeoxynucleotides. *J. Immunol.* 168: 4531–4537. [2385]
- Wu, C. *et al.* (2002) Distinct lineages of TH1 cells have differential capacities for memory cell generation *in vivo*. *Nat. Immunol.* 3: 852–858. [2680]
- Gutzmer, R. *et al.* (2003) Human Dendritic Cells Express the IL-18R and Are Chemoattracted to IL-18. *J. Immunol.* 171: 6363–6371. [4332]
- Holzer, U. *et al.* (2003) Differential Antigen Sensitivity and Costimulatory Requirements in Human Th1 and Th2 Antigen-Specific CD4⁺ Cells with Similar TCR Avidity. *J. Immunol.* 170: 1218–1223. [4348]
- Löhning, M. *et al.* (2003) Establishment of memory for IL-10 expression in developing T helper 2 cells requires repetitive IL-4 costimulation and does not impair proliferation. *PNAS* 100: 12307–12312. [3067]
- Schwaiger, S. *et al.* (2003) IL-4-Producing CD8⁺ T Cells with a CD62L^{++(bright)} Phenotype Accumulate in a Subgroup of Older Adults and Are Associated with the Maintenance of Intact Humoral Immunity in Old Age. *J. Immunol.* 170: 613–619. [4300]
- Voice, J.K. *et al.* (2003) Roles of Vasoactive Intestinal Peptide (VIP) in the Expression of Different Immune Phenotypes by Wild-Type Mice and T Cell-Targeted Type II VIP Receptor Transgenic Mice. *J. Immunol.* 170: 308–314. [4700]
- Guo, L. *et al.* (2004) Probabilistic Regulation of IL-4 Production in Th2 Cells: Accessibility at the *Il4* Locus. *Immunity* 20: 193–203. [4337]

- Li, L. *et al.* (2004) IL-4 Utilizes an Alternative Receptor to Drive Apoptosis of Th1 Cells and Skews Neonatal Immunity toward Th2. *Immunity* 20: 429–440. [4346]
- LeBlanc, R. *et al.* (2004) Immunomodulatory drug costimulates T cells via the B7-CD28 pathway. *Blood* 103: 1787–1790. [4344]
- Hata, H. *et al.* (2004) IL-18 together with anti-CD3 antibody induces human Th1 cells to produce Th1- and Th2-cytokines and IL-8. *Int. Immunol.* 16: 1733–1739. [4328]
- Tacchini-Cottier, F. *et al.* (2004) Notch1 expression on T cells is not required for CD4⁺ T helper differentiation. *Eur. J. Immunol.* 34: 1588–1596. [3325]
- Sojka, D.K. *et al.* (2004) IL-2 Secretion by CD4⁺ T Cells *In Vivo* Is Rapid, Transient, and Influenced by TCR-Specific Competition. *J. Immunol.* 172: 6136–6143. [4336]
- Ebert, L.M. *et al.* (2004) B cells alter the phenotype and function of follicular-homing CXCR5⁺ T cells. *Eur. J. Immunol.* 34: 3562–3571. [9016]
- Corbin, G.A. *et al.* (2005) T Cells Undergo Rapid ON/OFF but Not ON/OFF/ON Cycling of Cytokine Production in Response to Antigen. *J. Immunol.* 174: 718–726. [6613]
- McKarns, S.C. *et al.* (2005) Distinct Effects of TGF- β 1 on CD4⁺ and CD8⁺ T Cell Survival, Division, and IL-2 Production: A Role for T Cell Intrinsic Smad3. *J. Immunol.* 174: 2071–2083. [6766]
- Dannull, J. *et al.* (2005) Enhancing the immunostimulatory function of dendritic cells by transfection with mRNA encoding OX40 ligand. *Blood* 105: 3206–3213. [7045]
- Yang, J.Q. *et al.* (2003) Immunoregulatory Role of CD1d in the Hydrocarbon Oil-Induced Model of Lupus Nephritis. *J. Immunol.* 171: 2142–2153. [3097]
- Araki, M. *et al.* (2003) Th2 bias of CD4⁺ NKT cells derived from multiple sclerosis in remission. *Int. Immunol.* 15: 279–288. [4326]
- Yang, J.Q. *et al.* (2003) Repeated alpha-Galactosylceramide Administration Results in Expansion of NK T Cells and Alleviates Inflammatory Dermatitis in MRL-lpr/lpr Mice. *J. Immunol.* 171: 4439–4446. [4782]
- Desombre, I. *et al.* (2004) The interferon gamma secretion assay: a reliable tool to study interferon gamma production at the single cell level. *J. Immunol. Methods* 286: 167–185. [4323]
- Morris, S.C. *et al.* (2006) IL-4 Induces *In Vivo* Production of IFN- γ by NK and NKT Cells. *J. Immunol.* 176: 5299–5305. [8659]

NK cells

- Deniz, G. *et al.* (2002) Human NK1 and NK2 subsets determined by purification of IFN- γ -secreting and IFN- γ -nonsecreting NK cells. *Eur. J. Immunol.* 32: 879–884. [1279]
- Morris, S.C. *et al.* (2006) IL-4 Induces *In Vivo* Production of IFN- γ by NK and NKT Cells. *J. Immunol.* 176: 5299–5305. [8659]

DCs/ monocytes

- Edwards, A.D. *et al.* (2002) Microbial Recognition Via Toll-Like Receptor-Dependent and -Independent Pathways Determines the Cytokine Response of Murine Dendritic Cell Subsets to CD40 Triggering. *J. Immunol.* 169: 3652–3660. [2687]
- Fricke, I. *et al.* (2006) Mycobacteria Induce IFN- γ Production in Human Dendritic Cells via Triggering of TLR2. *J. Immunol.* 176: 5173–5182. [7059]

Regulatory T cells

- Veldman, C. *et al.* (2004) Type I Regulatory T Cells Specific for Desmoglein 3 Are More Frequently Detected in Healthy Individuals than in Patients with Pemphigus Vulgaris. *J. Immunol.* 172: 6468–6475. [5740]
- Akdis, M. *et al.* (2004) Immune Responses in Healthy and Allergic Individuals Are Characterized by a Fine Balance between Allergen-specific T Regulatory 1 and T Helper 2 Cells. *J. Exp. Med.* 199: 1567–1575. [3157]
- Sojka, D.K. *et al.* (2005) Early Kinetic Window of Target T Cell Susceptibility to CD25 Regulatory T Cell Activity. *J. Immunol.* 175: 7274–7280. [8169]

NKT cells

- Hayakawa, Y. *et al.* (2001) Critical Contribution of IFN- γ and NK cells, but not perforin-mediated cytotoxicity, to anti-metastatic effect of α -galactosylceramide. *Eur. J. Immunol.* 31: 1720–1727. [1073]
- Sfondrini, L. *et al.* (2002) Absence of the CD1 Molecule Up-Regulates Antitumor Activity Induced by CpG Oligodeoxynucleotides in Mice. *J. Immunol.* 169: 151–158. [2528]

Miltenyi Biotec



Miltenyi Biotec GmbH
Friedrich-Ebert-Straße 68
51429 Bergisch Gladbach
Germany
Phone +49 2204 8306-0
Fax +49 2204 85197
macs@miltenyibiotec.de

Miltenyi Biotec Inc.
12740 Earhart Avenue
Auburn CA 95602, USA
Phone 800 FOR MACS,
+1 530 888 8871
Fax +1 530 888 8925
macs@miltenyibiotec.com

Miltenyi Biotec Pty. Ltd.
(Australia)
Phone +61 02 8877 7400
macs@miltenyibiotec.com.au

Miltenyi Biotec B. V. (Benelux)
macs@miltenyibiotec.nl
Customer service, Netherlands
Phone 0800 4020120
Customer service, Belgium
Phone 0800 94016
Customer service, Luxembourg
Phone 800 24971

Miltenyi Biotec Shanghai Office
Phone +86 21 6235 1005
macs@miltenyibiotec.com.cn

Miltenyi Biotec (France)
Phone +33 1 56 98 16 16
macs@miltenyibiotec.fr

Miltenyi Biotec S.r.l. (Italy)
Phone +39 051 646 0411
macs@miltenyibiotec.it

Miltenyi Biotec K.K. (Japan)
Phone +81 3 56 46 8910
macs@miltenyibiotec.jp

Miltenyi Biotec Asia Pacific Pte. Ltd. (Singapore)
Phone +65 6238 8183
macs@miltenyibiotec.com.sg

Miltenyi Biotec S.L. (Spain)
Phone +34 91 512 12 90
macs@miltenyibiotec.es

Miltenyi Biotec Ltd. (UK)
Phone +44 1483 799 800
macs@miltenyibiotec.co.uk